

# Sparse Grid Quadrature and Interpolation

**Note:** We have encountered potentially high dimensional quadrature at several points:

$$\text{Bayes' Relation: } \pi(q|v_{obs}) = \frac{\pi(v_{obs}|q)\pi_0(q)}{\int_{\mathbb{R}^p} \pi(v_{obs}|q)\pi_0(q)dq}$$

$$\text{QoI: } y(t, x) = \mathbb{E}[u^K(t, x, Q)] = \int_{\Gamma} u^K(t, x, q)\rho_Q(q)dq$$

$$\text{Discrete Projection: } u_k(t, x) = \frac{1}{\gamma_k} \langle u, \Psi_k \rangle_{\rho} = \frac{1}{\gamma_k} \int_{\Gamma} u(t, x, q)\Psi_k(q)\rho_Q(q)dq$$

## Stochastic Quadrature Methods: Monte Carlo

$$\langle u, \Psi_k \rangle_{\rho} = \frac{1}{R} \sum_{r=1}^R u(t, x, q^r)\Psi_k(q^r)\rho_Q(q^r) + \varepsilon_R$$

### Notes:

- Errors satisfy  $\mathbb{E}[\varepsilon_R] = 0$  and  $\varepsilon_R = \mathcal{O}\left(\frac{1}{\sqrt{R}}\right)$  for large  $R$
- Optimal for sufficiently large  $p$ .

# Deterministic Quadrature Methods

## 1-D Quadrature Relations:

$$I^{(1)} f = \int_{\Gamma_1} f(q) \rho_Q(q) dq \approx \sum_{r=1}^R f(q^r) w^r = Q^{(1)} f$$

## Gaussian Quadrature:

$$I^{(1)} f = \frac{1}{2} \int_{-1}^1 f(q) dq \approx \frac{1}{2} \sum_{r=1}^R f(q^r) w^r,$$

$$I^{(1)} f = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} f(q) e^{-q^2/2} dq \approx \sum_{r=1}^R f(q^r) w^r$$

$r$	Nodes $q^r$	Weights $w^r$
1	0	2
2	$\pm \frac{1}{\sqrt{3}}$	1
3	0	$\frac{8}{9}$
	$\pm \sqrt{\frac{3}{5}}$	$\frac{5}{9}$
4	$\pm \frac{\sqrt{15+2\sqrt{30}}}{\sqrt{35}}$	$\frac{49}{6(18+\sqrt{30})}$
	$\pm \frac{\sqrt{15-2\sqrt{30}}}{\sqrt{35}}$	$\frac{49}{6(18-\sqrt{30})}$

# 1-D Quadrature Relations

**Nested Quadrature Techniques:** Consider on  $[0,1]$

$$Q_\ell^{(1)} f = \sum_{r=1}^{R_\ell} f(q_\ell^r) w_\ell^r$$

Trapezoid Rule:



$$Q_\ell^{(1)} f = \frac{1}{2h_\ell} \left[ f(0) + f(1) + 2 \sum_{r=1}^{R_\ell-1} f(q_\ell^r) \right]$$

where

$$h_\ell = \frac{1}{2^{\ell-1}}, \quad R_\ell = 2^{\ell-1} + 1, \quad q_\ell^r = rh_\ell = \frac{r}{2^{\ell-1}}, \quad w^r = \left[ \frac{1}{2h_\ell}, \frac{1}{h_\ell}, \dots, \frac{1}{h_\ell}, \frac{1}{2h_\ell} \right]$$

Clenshaw--Curtis:

$$q_\ell^r = \frac{1}{2} \left[ 1 - \cos \frac{\pi(r-1)}{R_\ell-1} \right], \quad r = 1, \dots, R_\ell$$



# Tensor Product Formulation

**Integrals:**

$$I^{(p)} f = \int_{\Gamma} f(q) \rho_Q(q) dq$$

**Tensor Product Quadrature:**

$$\begin{aligned} Q_{\ell}^{(p)} f &= \left( Q_{\ell_1}^{(1)} \otimes \cdots \otimes Q_{\ell_p}^{(1)} \right) f \\ &\equiv \sum_{r_1=1}^{R_{\ell_1}} \cdots \sum_{r_p=1}^{R_{\ell_p}} f(q_1^{r_1}, \dots, q_p^{r_p}) w_{\ell_1}^{r_1} \cdots w_{\ell_p}^{r_p} \end{aligned}$$

where


$$R = \prod_{i=1}^p R_{\ell_i}$$

**Errors:**

$$\left| I^{(p)} f - Q_{\ell}^{(p)} f \right| = \mathcal{O}(R_{\ell}^{-\alpha/p})$$

for functions  $f$  in the space

$$C^{\alpha}([0, 1]^p) = \left\{ f : [0, 1]^p \rightarrow \mathbb{R} \left| \max_{|\mathbf{k}'| \leq \alpha} \left\| \frac{\partial^{|\mathbf{k}'|} f}{\partial q_1^{k_1} \cdots \partial q_p^{k_p}} \right\|_{\infty} < \infty \right\}$$



# Sparse Grid Construction

## Motivation:

$R$							
0							1
1				$x$		$y$	
2			$x^2$		$xy$		$y^2$
3		$x^3$		$x^2y$		$xy^2$	$y^3$
4	$x^4$		$x^3y$		$x^2y^2$	$xy^3$	$y^4$

## Difference Relations: Define

$$\Delta_\ell^{(1)} f = \left( Q_\ell^{(1)} - Q_{\ell-1}^{(1)} \right) f$$

where

$$Q_\ell^{(1)} f = \sum_{r=1}^{R_\ell} f(q_\ell^r) w_\ell^r$$

### 1-D Nodal Points

$$\Theta_\ell^{(1)} = \left\{ q_\ell^1, \dots, q_\ell^{R_\ell} \right\}$$

## Example: Trapezoid rule

$$\ell = 2: \Theta_2^{(1)} = \left\{ 0, \frac{1}{2}, 1 \right\} \text{ and } w = \left[ \frac{1}{4}, \frac{1}{2}, \frac{1}{4} \right]$$

$$\ell = 1: \Theta_1^{(1)} = \{0, 1\} \text{ and } w = \left[ \frac{1}{2}, \frac{1}{2} \right]$$

Thus

$$\Delta_2^{(1)} f = -\frac{1}{4} f(0) + \frac{1}{2} f(1/2) - \frac{1}{4} f(1)$$

Note: Weights can be negative

# Sparse Grid Construction

## Sparse Grid Quadrature Rule:

$$\mathcal{Q}_\ell^{(p)} f = \sum_{|\ell'| \leq \ell + p - 1} \left( \Delta_{\ell_1}^{(1)} \otimes \cdots \otimes \Delta_{\ell_p}^{(1)} \right) f$$

where  $\ell' = (\ell_1, \dots, \ell_p) \in \mathbb{N}^p$  is a multi-index with  $|\ell'| = \sum_{i=1}^p \ell_i$ .

Note: Tensor product formula can be expressed as

$$\mathcal{Q}_\ell^{(p)} f = \sum_{\max \ell' \leq \ell} \left( \Delta_{\ell_1}^{(1)} \otimes \cdots \otimes \Delta_{\ell_p}^{(1)} \right) f$$

where  $\max \ell' \equiv \max\{\ell_1, \dots, \ell_p\}$ .




## Sparse Grid Nodal Set:

$$\Theta_\ell^{(p)} = \bigcup_{|\ell'| \leq \ell + p - 1} \Theta_{\ell_1}^{(1)} \times \cdots \times \Theta_{\ell_p}^{(1)}.$$

# Sparse Grid Construction

**Example:** Consider Clenshaw-Curtis with  $\Theta_1^{(1)} = \{\frac{1}{2}\}$  and  $\Theta_2^{(1)} = \{0, \frac{1}{2}, 1\}$ . For  $p = 2$ ,  $\ell' = (\ell_1, \ell_2)$  so  $|\ell'| = \ell_1 + \ell_2$ . For  $\ell = 4$ , the sparse grid nodal set is

$$\begin{aligned} \Theta_4^{(2)} &= \left( \Theta_1^{(1)} \times \Theta_1^{(1)} \right) \cup \left( \Theta_1^{(1)} \times \Theta_2^{(1)} \right) \cup \left( \Theta_2^{(1)} \times \Theta_1^{(1)} \right) \\ &\cup \left( \Theta_1^{(1)} \times \Theta_3^{(1)} \right) \cup \left( \Theta_2^{(1)} \times \Theta_2^{(1)} \right) \cup \left( \Theta_3^{(1)} \times \Theta_1^{(1)} \right) \\ &\cup \left( \Theta_1^{(1)} \times \Theta_4^{(1)} \right) \cup \left( \Theta_2^{(1)} \times \Theta_3^{(1)} \right) \cup \left( \Theta_3^{(1)} \times \Theta_2^{(1)} \right) \cup \left( \Theta_4^{(1)} \times \Theta_1^{(1)} \right) \end{aligned}$$



Sparse Grid: 29 points

Tensored Grid: 81 points

# Sparse Grid Construction

## Error Analysis:

$$\|\mathcal{I}f - \mathcal{A}(q, p)f\| = \mathcal{O}\left(\mathcal{R}^{-\alpha} \log(\mathcal{R})^{(p-1)(\alpha+1)}\right)$$

## Grid Sizes:

$p$	$R_\ell$	Sparse Grid $\mathcal{R}$	Tensor Grid $R = (R_\ell)^p$
2	5	13	25
	9	29	81
5	5	61	3125
	9	241	59,049
10	5	221	9,765,625
	9	1581	$> 3 \times 10^9$
50	5	5101	$> 8 \times 10^{34}$
	9	171,901	$> 5 \times 10^{47}$
100	5	20,201	$> 7 \times 10^{69}$
	9	1,353,801	$> 2 \times 10^{95}$





# Anisotropic Sparse Grids

**Multi-Index Set:**

$$\mathbb{I}(\ell) = \left\{ \ell' \in \mathbb{N}^p \mid \ell' \cdot \mathbf{a} = \sum_{i=1}^p a_i \ell'_i \leq \ell + p - 1 \right\}$$

where  $\mathbf{a} \in \mathbb{R}_+^p$  is a vector of weights.



# Interpolating Polynomials

**1-D Interpolation:** Seek polynomials that satisfy

$$u^{M_1}(q^m) = u^m = u(q^m), \quad m = 1, \dots, M_1$$

**Lagrange Interpolation:** Employ representation

$$u^{M_1}(q) = \sum_{m=1}^{M_1} u^m L_m(q)$$

where  $L_m(q)$  are Lagrange interpolating polynomials defined by

$$\begin{aligned} L_m(q) &= \prod_{\substack{j=0 \\ j \neq m}}^{M_1} \frac{q - q^j}{q^m - q^j} \\ &= \frac{(q - q^1) \cdots (q - q^{m-1})(q - q^{m+1}) \cdots (q - q^{M_1})}{(q^m - q^1) \cdots (q^m - q^{m-1})(q^m - q^{m+1}) \cdots (q^m - q^{M_1})} \end{aligned}$$

**Note:** By construction, the Lagrange polynomials satisfy





$$L_m(q^n) = \delta_{mn}, \quad 1 \leq m, n \leq M_1$$

# 1-D Interpolating Polynomials

**Example:** Consider the Runge function  $f(q) = \frac{1}{1 + 25q^2}$  with points

$$q^j = -1 + (j - 1) \frac{2}{M_1}, \quad j = 1, \dots, M_1 + 1$$

$$q^j = -\cos \frac{\pi(j - 1)}{M_1 - 1}, \quad j = 1, \dots, M_1$$



# Multi-Dimensional Interpolation

**Tensor Products and Sparse Grid:** Interpolation formulae analogous to quadrature rules.

## **Uses:**

- Spectral collocation methods to propagate input uncertainties.
- Construction of surrogate models.

## **Sparse Grid Software:**

- MATLAB: Sparse Grid Interpolation Toolbox – Be careful of Clenshaw-Curtis, which are actually Newton-Cotes points – versus Chebychev
- Dakota